

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : C12N 15/82, 15/55, 9/16 C12N 1/21, A23K 1/165 A01H 5/00	A1	(11) International Publication Number: WO 91/14782 (43) International Publication Date: 3 October 1991 (03.10.91)
(21) International Application Number: PCT/NL91/00047 (22) International Filing Date: 25 March 1991 (25.03.91) (30) Priority data: 498,561 23 March 1990 (23.03.90) US 586,765 21 September 1990 (21.09.90) US (71) Applicants: GIST-BROCADES N.V. [NL/NL]; Wateringseweg 1, P.O. Box 1, NL-2600 MA Delft (NL). MOGEN INTERNATIONAL N.V. [NL/NL]; Einsteinweg 97, NL-2333 CB Leiden (NL). (72) Inventors: PEN, Jan ; Voszegge 9, NL-2318 ZJ Leiden (NL). HOEKEMA, Andreas ; Warmonderweg 66, NL-2341 KX Oegstgeest (NL). SIJMONS, Peter, Christiaan ; Valeriusstraat 210-3, NL-1075 GK Amsterdam (NL). VAN OOYEN, Albert, J., J. ; Overburgkade 78, NL-2275 GK Amsterdam (NL). RIETVELD, Krijn ; Van der Walstraat 129, NL-3132 TL Vlaardingen (NL). VERWOERD, Teunis, Cornelis ; Oudesingel 286, NL-2312 RK Leiden (NL).		(74) Agents: HUYGENS, Arthur, Victor et al.; Gist-Brocades N.V., Patents and Trademarks Department, Wateringseweg 1, PO-box 1, NL-2600 MA Delft (NL). (81) Designated States: AU, CA, FI, HU, JP, KR, SU. Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: THE EXPRESSION OF PHYTASE IN PLANTS (57) Abstract <p>The present invention provides for the expression of phytase in transgenic plants or plant organs and methods for the production of such plants. DNA expression constructs are provided for the transformation of plants with a gene encoding phytase under the control of regulatory sequences which are capable of directing the expression of phytase. These regulatory sequences include sequences capable of directing transcription in plants, either constitutively, or stage and/or tissue specific, depending on the use of the plant or parts thereof. The transgenic plants and plant organs provided by the present invention may be applied to a variety of industrial processes either directly, e.g. in animal feeds or alternatively, the expressed phytase may be extracted and if desired, purified before application.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

5

THE EXPRESSION OF PHYTASE IN PLANTS

Field of the Invention

10 The present invention pertains to the production of phytase in transgenic plants and the use of the thus-produced phytase in industrial processes.

Background of the Invention

15 Phosphorus is an essential element for the growth of all organisms. In livestock production, feed must be supplemented with inorganic phosphorus in order to obtain a good growth performance of monogastric animals (e.g. pigs, poultry and fish).

20 In contrast, no inorganic phosphate needs to be added to the feedstuffs of ruminant animals. Microorganisms, present in the rumen, produce enzymes which catalyze the conversion of phytate (myo-inositolhexakis-phosphate) to inositol and inorganic phosphate.

25 Phytate occurs as a storage phosphorus source in virtually all feed substances originating from plants (for a review see: Phytic acid, chemistry and applications, E. Graf (ed.), Pilatus Press; Minneapolis, MN, U.S.A. (1986)). Phytate comprises 1-3% of all nuts, cereals, legumes, oil seeds, spores and pollen. Complex salts of phytic acid are termed phytin. Phytic acid is considered to be an anti-
30 nutritional factor since it chelates minerals such as calcium, zinc, magnesium, iron and may also react with proteins, thereby decreasing the bioavailability of proteins and nutritionally important minerals.

35 Phytate phosphorus passes through the gastrointestinal tract of monogastric animals and is excreted in the manure. Though some hydrolysis of phytate does occur in the colon, the thus-released inorganic phosphorus has no nutritional value since inorganic phosphorus is absorbed

only in the small intestine. As a consequence, a significant amount of the nutritionally important phosphorus is not used by monogastric animals, despite its presence in the feed.

5 The excretion of phytate phosphorus in manure has further consequences. Intensive livestock production has increased enormously during the past decades. Consequently, the amount of manure produced has increased correspondingly and has caused environmental problems in various parts of the world. This is due, in part, to the accumulation of
10 phosphate from manure in surface waters which has caused eutrophication.

The enzymes produced by microorganisms, which catalyze the conversion of phytate to inositol and inorganic phosphorus are broadly known as phytases. Phytase producing
15 microorganisms comprise bacteria such as Bacillus subtilis (V.K. Paver and V.J. Jagannathan (1982) J. Bacteriol. 151, 1102) and Pseudomonas (D.J. Cosgrove (1970) Austral. J. Biol. Sci. 23, 1207); yeasts such as Saccharomyces cerevisiae (N.R. Nayini and P. Markakis (1984) Lebensmittel
20 Wissenschaft und Technologie 17, 24); and fungi such as Aspergillus terreus (K. Yamada, Y. Minoda and S. Yamamoto (1986) Agric. Biol. Chem. 32, 1275). Various other Aspergillus species are known to produce phytase, of which, the phytase produced by Aspergillus ficuum has been
25 determined to possess one of the highest levels of specific activity, as well as having better thermostability than phytases produced by other microorganisms (van Gorcom et al. (1991) European Patent Application 89202436.5, Publication No. 0 420 358, filed September 27, 1989).

30 Phytases are also endogenously present in many plant species (see Loewus, F.A. (1990) In: Plant Biology vol. 9: "Inositol metabolism in plants" (eds. D.J. Morré, W.F. Boss, F.A. Loewus) 13). Gellatly, K.S. and Lefebvre, D.D. ((1990) Plant Physiology (supplement), 93, abstract 562) mention the
35 isolation and characterization of a phytase cDNA clone obtained from potato tubers. Gibson, D.M. et al. and

Christen, A.A. et al. ((1988) J. Cell Biochem., 12C,
abstracts L407 and L402, respectively) mention the synthesis
of endogenous phytase during the germination of soybean
seeds. However, plant phytases are normally produced in
5 amounts insufficient for their application in industrial
processes, per se.

The concept of adding microbial phytase to the
feedstuffs of monogastric animals has been previously
described (Ware, J.H., Bluff, L. and Shieh, T.R. (1967) U.S.
10 Patent No. 3,297,548; Nelson, T.S., Shieh, T.R., Wodzinski,
R.J. and Ware, J.H. (1971) J. Nutrition 101, 1289). To date,
however, application of this concept has not been
commercially feasible, due to the high cost of the
production of the microbial enzymes (Y.W. Han (1989) Animal
15 Feed Sci. and Technol. 24, 345). For economic reasons,
inorganic phosphorus is still added to monogastric animal
feedstuffs.

Phytases have found other industrial uses as well.
Exemplary of such utilities is an industrial process for the
20 production of starch from cereals such as corn and wheat.
Waste products comprising e.g. corn gluten feeds from such a
wet milling process are sold as animal feed. During the
steeping process phytase may be supplemented. Conditions
($T \approx 50^{\circ}\text{C}$ and $\text{pH} = 5.5$) are ideal for fungal phytases (see
25 e.g. European Patent Application 0 321 004 to Alko Ltd.).
Advantageously, animal feeds derived from the waste products
of this process will contain phosphate instead of phytate.

It has also been conceived that phytases may be used
in soy processing (see FinaseTM Enzymes By Alko, a product
30 information brochure published by Alko Ltd., Rajamäki,
Finland). Soybean meal contains high levels of the anti-
nutritional factor phytate which renders this protein source
unsuitable for application in baby food and feed for fish,
calves and other non-ruminants. Enzymatic upgrading of this
35 valuable protein source improves the nutritional and
commercial value of this material.

The possibility of using transgenic plants as a production system for valuable proteins has been proposed. Examples to date are the production of interferon in tobacco (Goodman, R.M., Knauf, V.C., Houck, C.M. and Comai, L. (1987) PCT/WO 87/00865), enkephalins in tobacco, Brassica napus and Arabidopsis thaliana (Vandekerckhove, J., Van Damme, J., Van Lijsebettens, M., Botterman, J., DeBlock, M., DeClerq, A., Leemans, J., Van Montagu, M. and Krebbers, E. (1989) Bio/Technol. 7, 929), antibodies in tobacco (Hiatt, A., Cafferkey, R. and Boedish, K. (1990) Nature 342, 76) and human serum albumin in tobacco and potato (Sijmons, P.C., Dekker, B.M.M., Schrammeijer, B., Verwoerd, T.C., van den Elzen, P.J.M. and Hoekema, A. (1990) Bio/Technol. 8, 217).

In practice, the transformation of an increasing number of plant species, especially dicotyledonous species (e.g. tobacco, potato, tomato, Petunia, Brassica), has become a routine procedure for workers skilled in the art (Klee, H., Horsch, R. and Rogers, S. (1987) Annu. Rev. Plant Physiol. 38, 467; Gasser C.S. and Fraley, R.T. (1989) Science 244, 1293). Strategies for the expression of foreign genes in plants have become well established (Gasser and Fraley, supra). Regulatory sequences from plant genes have been identified that are used for the construction of chimeric genes that can be functionally expressed in plants and plant cells.

For the introduction of gene constructions into plants, several technologies are available, such as transformation with Agrobacterium tumefaciens or Agrobacterium rhizogenes. Using this strategy, a wide variety of plant tissues have been exploited, the choice being largely dependent on the plant species and its amenability in tissue culture. Successful examples are the transformation of protoplasts, microspores or pollen, and explants such as leaves, stems, roots, hypocotyls and cotyls. Furthermore, methods for direct DNA introduction in protoplasts and plant cells or tissues are used such as

microinjection, electroporation, particle bombardment and direct DNA uptake (Gasser and Fraley, supra).

Proteins may be produced in plants using a variety of expression systems. For instance, the use of a constitutive promoter such as the 35S promoter of Cauliflower Mosaic Virus (CaMV) (Guilley, H., Dudley, R.K., Jonard, G., Balazs, E. and Richards, K.E. (1982) Cell 30, 763) will result in the accumulation of the expressed protein in all organs of the transgenic plant. Alternatively, use may be made of promoters from genes encoding proteins which are expressed in a highly tissue-specific and stage-specific manner (Higgins, T.J.V., (1984) Annu. Rev. Plant Physiol. 35, 191; Shotwell, M.A. and Larkins, B.A. (1989) In: The biochemistry of plants Vol. 15 (Academic Press, San Diego: Stumpf, P.K. and Conn, E.E., eds.), 297), i.e., the genes are expressed only in the target tissue and only during the desired stage of development.

It will be appreciated that an economical procedure for the production of phytase will be of significant benefit to, inter alia, the animal feed industry. One method of producing a more economical phytase would be to use recombinant DNA techniques to produce transgenic plants or plant organs capable of expressing phytase which could then in turn be added as such, for example, to animal food or feedstuffs for direct consumption by the animal. Alternatively, the phytase expressed in these transgenic plants or plant organs could be extracted and if desired, purified for the desired application.

30 Summary of the Invention

The present invention provides for the expression of phytase in transgenic plants or plant organs and methods for the production of such plants. This is achieved via the introduction into the plant of an expression construct comprising a DNA sequence encoding a protein having phytase activity.

The DNA expression constructs provided by the present invention for the transformation of plants are under the control of regulatory sequences which are capable of directing the expression of phytase. These regulatory sequences may also include sequences capable of directing transcription in plants, either constitutively, or stage and/or tissue specific, depending on the use of the plant or parts thereof.

The transgenic plants and plant organs provided by the present invention may be applied to a variety of industrial processes either directly, e.g. in animal food or feedstuffs or alternatively, the expressed phytase may be extracted and if desired, purified before application.

Brief Description of the Figures

- Figure 1. Strategy for the cloning of phytase cDNA.
- Figure 2. Nucleotide sequence of the translated region of the phytase cDNA fragment and the derived amino acid sequence of the phytase protein; the start of the mature phytase protein is indicated as position +1.
- Figure 3. Binary vector pMOG23.
- Figure 4. Oligonucleotide duplexes used in cloning.
- Figure 5. Plasmid pMOG29. Plasmid pUC18 containing an expression cassette for constitutive expression in plants and a sequence encoding a tobacco signal peptide.
- Figure 6. The effects of the addition of ground seeds containing phytase on the liberation of inorganic phosphorous from phytate.
- Figure 7. Dose-response relationship of Aspergillus phytase in an in vitro digestion model.
- Figure 8. Dose-response relationship of Aspergillus phytase and phytase contained in tobacco seed in an in vitro digestion model.

Detailed Description of the Invention

According to the present invention, transgenic plants or plant organs are obtained in which phytase is produced. This is achieved via the introduction into the plant of an expression construct comprising a DNA sequence encoding a protein having phytase activity.

DNA expression constructs are provided by the present invention for the stable transformation of plants with a gene encoding a phytase. These constructs comprise a DNA sequence encoding a phytase which is operably linked to regulatory sequences which are capable of directing the expression of phytase. These regulatory sequences may also include sequences capable of directing transcription in plants, either constitutively, or stage and/or tissue specific, depending on the use of the plant or parts thereof.

The expression constructs provided by the present invention may be inserted into a vector, preferably a plasmid, used in bacteria-mediated transformation of the selected plant host. The expression construct is then preferably integrated into the genome of the plant host.

Within the context of the present invention, the term phytase embraces a family of enzymes which catalyze reactions involving the liberation of inorganic phosphorus from various myoinositol phosphates. This is understood to embrace all proteins having phytase activity.

The DNA sequence encoding phytase may be obtained from a variety of sources such as microbial, plant or animal sources. Preferably, the DNA sequence is obtained from a microbial source such as the filamentous fungus Aspergillus. Most preferred DNA sequences are obtained from Aspergillus ficuum, Aspergillus niger, Aspergillus awamori and Aspergillus nidulans.

The cloning of a gene or a cDNA encoding a phytase protein may be achieved using various methods. One method is by purification of the phytase protein, subsequent

determination of the N-terminal and several internal amino acid sequences and screening of a genomic or cDNA library of the organism producing the phytase, using oligonucleotide probes based on the amino acid sequences.

5 If at least a partial sequence of the gene is known, this information may be used to clone the corresponding cDNA using, for instance, the polymerase chain reaction (PCR) (PCR Technology: Principles and Applications for DNA Amplification, (1989) H.A. Ehrlich, ed., Stockton Press, New York).

10 It will be evident to those skilled in the art that the cloned phytase gene described above may be used in heterologous hybridization experiments, directed to the isolation of phytase encoding genes from other micro-organisms.

15 In another aspect, the cloned phytase gene described above may be used as starting materials for the construction of "second generation" phytases. "Second generation" phytases are phytases, altered by mutagenesis techniques (e.g. site-directed mutagenesis), which have properties that differ from those of wild-type phytases or recombinant phytases such as those produced by the present invention. For example, the temperature or pH optimum, specific activity or substrate affinity may be altered so as to be better suited for application in a defined process.

20 The isolation of the cDNA encoding phytase enables the construction of expression constructs capable of directing the production of phytase in the selected plant host via the application of recombinant DNA techniques such as the exchange of regulatory elements such as e.g. promoters, secretional signals, or combinations thereof.

30 Phytase may be produced constitutively in the transgenic plants during all stages of development. Depending on the use of the plant or plant organs, the enzymes may be expressed in a stage-specific manner, for instance during tuber formation or fruit development. Also,

depending on the use, the enzymes may be expressed tissue-specifically, for instance in plant organs such as fruit, tubers, leaves or seeds.

5 Transgenic plants, as defined in the context of the present invention include plants (as well as parts and cells of said plants) and their progeny, which have been genetically modified using recombinant DNA techniques to cause or enhance production of a phytase in the desired plant or plant organ.

10 In the context of the present invention, the phrase "an enhanced amount of phytase" refers specifically to a statistically significant amount of plant tissue which, on average, contain a statistically significant greater amount of phytase as compared with the average amount of phytase
15 enzyme found in an equal amount of non-modified plant tissue.

Within the context of the present invention, plants to be selected include, but are not limited to crops producing edible flowers such as cauliflower (Brassica oleracea),
20 artichoke (Cynara scolymus), fruits such as apple (Malus, e.g. domesticus), banana (Musa, e.g. acuminata), berries (such as the currant, Ribes, e.g. rubrum), cherries (such as the sweet cherry, Prunus, e.g. avium), cucumber (Cucumis, e.g. sativus), grape (Vitis, e.g. vinifera), lemon (Citrus
25 limon), melon (Cucumis melo), nuts (such as the walnut, Juglans, e.g. regia; peanut, Arachis hypogaeae), orange (Citrus, e.g. maxima), peach (Prunus, e.g. persica), pear (Pyra, e.g. communis), plum (Prunus, e.g. domestica), strawberry (Fragaria, e.g. moschata), tomato (Lycopersicon,
30 e.g. esculentum), leafs, such as alfalfa (Medicago, e.g. sativa), cabbages (e.g. Brassica oleracea), endive (Cichoreum, e.g. endivia), leek (Allium, e.g. porrum), lettuce (Lactuca, e.g. sativa), spinach (Spinacia e.g. oleraceae), tobacco (Nicotiana, e.g. tabacum), roots, such
35 as arrowroot (Maranta, e.g. arundinacea), beet (Beta, e.g. vulgaris), carrot (Daucus, e.g. carota), cassava (Manihot,

e.g. esculenta); turnip (Brassica, e.g. rapa), radish (Raphanus, e.g. sativus), yam (Dioscorea, e.g. esculenta), sweet potato (Ipomoea batatas) and seeds, such as bean (Phaseolus, e.g. vulgaris), pea (Pisum, e.g. sativum), soybean (Glycin, e.g. max), wheat (Triticum, e.g. aestivum), barley (Hordeum, e.g. vulgare), corn (Zea, e.g. mays), rice (Oryza, e.g. sativa), rapeseed (Brassica napus), millet (Panicum L.), sunflower (Helianthus annus), oats (Avena sativa), tubers, such as kohlrabi (Brassica, e.g. oleraceae), potato (Solanum, e.g. tuberosum) and the like.

The choice of the plant species is primarily determined by the intended use of the plant or parts thereof and the amenability of the plant species to transformation.

Several techniques are available for the introduction of the expression construct containing the phytase-encoding DNA sequence into the target plants. Such techniques include but are not limited to transformation of protoplasts using the calcium/polyethylene glycol method, electroporation and microinjection or (coated) particle bombardment (Potrykus, I. (1990) Bio/Technol. 8, 535).

In addition to these so-called direct DNA transformation methods, transformation systems involving vectors are widely available, such as viral vectors (e.g. from the Cauliflower Mosaic Virus (CaMV) and bacterial vectors (e.g. from the genus Agrobacterium) (Potrykus, supra). After selection and/or screening, the protoplasts, cells or plant parts that have been transformed can be regenerated into whole plants, using methods known in the art (Horsch, R.B., Fry, J.E., Hoffmann, N.L., Eichholtz, D., Rogers, S.G. & Fraley, R.T. (1985) Science 227, 1229). The choice of the transformation and/or regeneration techniques is not critical for this invention.

For dicots, a preferred embodiment of the present invention uses the principle of the binary vector system (Hoekema, A., Hirsch, P.R., Hooykaas, P.J.J. and Schilperoort, R.A. (1983) Nature 303, 179; Schilperoort,

R.A., Hoekema, A. and Hooykaas, P.J.J. (1984) European Patent Application No. 0 120 516) in which Agrobacterium strains are used which contain a vir plasmid with the virulence genes and a compatible plasmid containing the gene construct to be transferred. This vector can replicate in both E. coli and in Agrobacterium, and is derived from the binary vector Bin19 (Bevan, M. (1984) Nucl. Acids Res. 12, 8711) which is altered in details that are not relevant for this invention. The binary vectors as used in this example contain between the left- and right-border sequences of the T-DNA, an identical NPTII-gene coding for kanamycin resistance (Bevan, supra) and a multiple cloning site to clone in the required gene constructs.

The transformation and regeneration of monocotyledonous crops is not a standard procedure. However, recent scientific progress shows that in principle monocots are amenable to transformation and that fertile transgenic plants can be regenerated from transformed cells. The development of reproducible tissue culture systems for these crops, together with the powerful methods for introduction of genetic material into plant cells has facilitated transformation. Presently the methods of choice for transformation of monocots are microprojectile bombardment of explants or suspension cells, and direct DNA uptake or electroporation of protoplasts. For example, transgenic rice plants have been successfully obtained using the bacterial hph gene, encoding hygromycin resistance, as a selection marker. The gene was introduced by electroporation (Shimamoto, K., Terada, R., Izawa, T. and Fujimoto, H. (1989) Nature 338, 274). Transgenic maize plants have been obtained by introducing the Streptomyces hygrosopicus bar gene, which encodes phosphinothricin acetyltransferase (an enzyme which inactivates the herbicide phosphinothricin), into embryogenic cells of a maize suspension culture by microparticle bombardment (Gordon-Kamm, W.J., Spencer, T.M., Mangano, M.L., Adams, T.R., Daines, R.J., Start, W.G.,

O'Brien, J.V., Chambers, S.A., Adams Jr., W.R., Willets, N.G., Rice, T.B., Mackey, C.J., Krueger, R.W., Kausch, A.P. and Lemaux, P.G. (1990) *The Plant Cell* 2, 603). The introduction of genetic material into aleurone protoplasts of other monocot crops such as wheat and barley has been reported (Lee, B., Murdoch, K., Topping, J., Kreis, M. and Jones, M.G.K. (1989) *Plant Mol. Biol.* 13, 21). Wheat plants have been regenerated from embryogenic suspension culture by selecting only the aged compact and nodular embryogenic callus tissues for the establishment of the embryogenic suspension cultures (Vasil, V., Redway, F. and Vasil, I.K. (1990) *Bio/Technol.* 8, 429). The combination with transformation systems for these crops enables the application of the present invention to monocots. These methods may also be applied for the transformation and regeneration of dicots.

Expression of the phytase construct involves such details as transcription of the gene by plant polymerases, translation of mRNA, etc. that are known to persons skilled in the art of recombinant DNA techniques. Only details relevant for the proper understanding of this invention are discussed below.

Regulatory sequences which are known or are found to cause expression of phytase may be used in the present invention. The choice of the regulatory sequences used depends on the target crop and/or target organ of interest. Such regulatory sequences may be obtained from plants or plant viruses, or may be chemically synthesized. Such regulatory sequences are promoters active in directing transcription in plants, either constitutively or stage and/or tissue specific, depending on the use of the plant or parts thereof. These promoters include, but are not limited to promoters showing constitutive expression, such as the 35S promoter of Cauliflower Mosaic Virus (CaMV) (Guilley et al. (1982) *Cell* 30, 763), those for leaf-specific expression, such as the promoter of the ribulose

bisphosphate carboxylase small subunit gene (Coruzzi et al., (1984) EMBO J. 3, 1671), those for root-specific expression, such as the promoter from the glutamine synthase gene (Tingey et al. (1987) EMBO J., 6, 3565), those for seed-specific expression, such as the cruciferin A promoter from Brassica napus (Ryan et al. (1989) Nucl. Acids Res. 17, 3584), those for tuber-specific expression, such as the class-I patatin promoter from potato (Rocha-Sosa et al., (1989) EMBO J. 8, 23; Wenzler et al., (1989) Plant Mol. Biol. 12, 41) or those for fruit-specific expression, such as the polygalacturonase (PG) promoter from tomato (Bird et al., (1988) Plant Mol. Biol. 11, 651).

Other regulatory sequences such as terminator sequences and polyadenylation signals include any such sequence functioning as such in plants, the choice of which is within the level of the skilled artisan. An example of such sequences is the 3' flanking region of the nopaline synthase (nos) gene of Agrobacterium tumefaciens (Bevan, M., supra).

The regulatory sequences may also include enhancer sequences, such as found in the 35S promoter of CaMV, and mRNA stabilizing sequences such as the leader sequence of Alfalfa Mosaic Virus (AlMV) RNA4 (Brederode, F.T., Koper-Zwarthoff, E.C. & Bol, J.F. (1980) Nucl. Acids Res. 8, 2213) or any other sequences functioning in a like manner.

The phytase should be expressed in an environment that allows for stability of the expressed protein. The choice of cellular compartments, such as cytosol, endoplasmic reticulum, vacuole, protein body or periplasmic space can be used in the present invention to create such a stable environment, depending on the biophysical parameters of the phytase. Such parameters include, but are not limited to pH-optimum, sensitivity to proteases or sensitivity to the molarity of the preferred compartment.

To obtain expression in the cytoplasm of the cell, the expressed enzyme should not contain a secretory signal

peptide or any other target sequence. For expression in chloroplasts and mitochondria the expressed enzyme should contain a specific so-called transit peptide for import into these organelles. Targeting sequences that can be attached to the enzyme of interest in order to achieve this are known (Smeekens et al, (1990) T.I.B.S. 15, p.73; van den Broeck et al., (1985) Nature 313, 358; Schreier et al., (1985) EMBO J. 4, 25). If the activity of the enzyme is desired in the vacuoles a secretory signal peptide has to be present, as well as a specific targeting sequence that directs the enzyme to these vacuoles (Tague et al, (1988) Plant Phys. 86, 506). The same is true for the protein bodies in seeds. The DNA sequence encoding the enzyme of interest should be modified in such a way that the enzyme can exert its action at the desired location in the cell.

To achieve extracellular expression of the phytase, the expression construct of the present invention utilizes a secretory signal sequence. Although signal sequences which are homologous (native) to the plant host species are preferred, heterologous signal sequences, i.e. those originating from other plant species or of microbial origin, may be used as well. Such signal sequences are known to those skilled in the art. Appropriate signal sequences which may be used within the context of the present invention are disclosed in Walter, P. and Blobel, G. (1986) Biochem. Soc. Symp., 47, 183; Von Heijne, G. (1986) J. Mol. Biol., 189, 239; and Sijmons, P.C., Dekker, B.M.M., Schrammeijer, B., Verwoerd, T.C., van den Elzen, P.J.M. and Hoekema, A. (1990) Bio/Technol., 8, 217.

All parts of the relevant DNA constructs (promoters, regulatory-, secretory-, stabilizing-, targeting- or termination sequences) of the present invention may be modified, if desired, to affect their control characteristics using methods known to those skilled in the art.

It is pointed out that plants containing phytase obtained via the present invention may be used to obtain plants or plant organs with yet higher phytase levels. For example, it may be possible to obtain such plants or plant organs by the use of somoclonal variation techniques or by cross breeding techniques. Such techniques are well known to those skilled in the art.

In one embodiment of the present invention, a double-stranded cDNA encoding phytase is prepared from mRNA isolated from Aspergillus ficuum. The DNA construct is placed under the control of regulatory sequences from the gene encoding the 12S storage protein cruciferin from Brassica napus. The construct is thereafter subcloned into a binary vector such as pMOG23 (in E. coli K-12 strain DH5 α , deposited at the Centraal Bureau voor Schimmelcultures, Baarn, The Netherlands on January 29, 1990 under accession number CBS 102.90). This vector is introduced into Agrobacterium tumefaciens which contains a disarmed Ti plasmid. Bacterial cells containing this construct are co-cultivated with tissues from tobacco or Brassica plants, and transformed plant cells are selected by nutrient media containing antibiotics and induced to regenerate into differentiated plants on such media. The resulting plants will produce seeds that contain and express the DNA construct.

In another embodiment of the present invention, the phytase-encoding DNA construct is placed under the control of regulatory sequences from the 35S promoter of Cauliflower Mosaic Virus (CaMV). The construct is thereafter subcloned into a binary vector. This vector is then introduced into Agrobacterium tumefaciens which contains a disarmed Ti plasmid. Bacterial cells containing this construct are co-cultivated with tissues from tobacco or Brassica plants, and transformed plant cells are selected by nutrient media containing antibiotics and induced to regenerate into

differentiated plants on such media. The resulting plants contain and express the DNA construct constitutively.

Phytase activity may be measured via a number of assays, the choice of which is not critical to the present invention. For example, the phytase enzyme activity of the transgenic plant tissue may be tested with an ELISA-assay, Western blotting or direct enzyme assays using colorimetric techniques or native gel assays.

The plant or plant organ containing phytase, as produced via the present invention may be used in a variety of industrial processes requiring the action of a phytase.

The plants or plant organs containing phytase produced according to the present invention may be used in industrial processes requiring the action of a phytase. Examples of such applications are in feed additives for non-ruminants, in soy processing, or in the production of inositol or inositol-phosphates from phytate. Other industrial processes using substrates that contain phytate such as the starch industry and in fermentation industries, such as the brewing industry. Chelation of metal ions by phytate may cause these minerals to be unavailable for the production microorganisms. Enzymatic hydrolysis of phytate prevents these problems.

Phytase produced in plants can also be used in a process for steeping corn or sorghum kernels. The plant tissue may be ground before adding to steeping corn. Phytase liberated from the plant tissue can act on phytin, which is present in many corn preparations. Degradation of phytin in steeping corn is beneficial for the added commercial value of corn steep liquor, which is used as animal feed or as a nutrient in microbial fermentations. Furthermore, the degradation of phytin can prevent problems relating to the accumulation of deposits in filters, pipes, reactor vessels, etc. during concentration, transport and storage of corn steep liquor (Vaara, T. et al. (1989) European Patent Application 0 321 004). The action of phytase can also

accelerate the steeping process and the separation processes involved in corn wet milling.

5 The plants or plant organs may be used directly, i.e. without further processing, or may first be processed via conventional means such as grinding to the desired consistency before application.

10 Alternatively, the phytase may be extracted from the plant or plant organ and, if desired, purified before use using conventional extraction methods and purification techniques.

15 The production of phytases in plants which are compatible with the intended application provides convenience and will reduce production costs as compared to that of microbial phytases in order to allow its economical application, e.g. in animal feed, which eventually will lead to a price/in vivo performance ratio competitive with inorganic phosphate. As a further benefit, the phosphorus content of manure will be considerably decreased.

20 It will be appreciated that the application of phytases, available at a price competitive with inorganic phosphate, will increase the degrees of freedom for the compound feed industry to produce a high quality feed. For example, when feed is supplemented with phytase, the addition of inorganic phosphate may be omitted and the contents of various materials containing phytate may be increased.

30 The following examples are provided so as to give those of ordinary skill in the art a complete disclosure and description of how to make and use the invention and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, pH, etc.) but some experimental errors and deviation should be accounted for. Unless indicated otherwise, temperature is in degrees Centigrade and pressure is at or near atmospheric.

Example 1

Isolation of poly A⁺ RNA from Aspergillus ficuum

5 A. ficuum strain NRRL 3135 is grown in a medium containing 22.72 g/l maize flour (amylase treated at pH 7 at 85°C during 15 minutes), 9.36 g/l glucose, 2.9 g/l KNO₃, 0.142 g/l KCl, 0.142 g/l MgSO₄·7H₂O and 56.8 mg/l FeSO₄·7H₂O. After 6 days the mycelium is harvested.

10 Dry mycelium (0.5 g) is frozen with liquid nitrogen and ground. Subsequently the material is homogenized with an Ultra turrax (full speed, 1 minute) at 0°C in 3 M LiCl, 6 M Urea and maintained overnight at 4°C as described by Auffray and Rougeon (1980) Eur. J. Biochem. 107, 303. Total cellular RNA is obtained after centrifugation at 16,000 X g, which is
15 followed by two successive extractions with phenol:chloroform:isoamylalcohol (50:48:2). The RNA is precipitated with ethanol and redissolved in 1 ml 10 mM Tris-HCl (pH 7.4), 0.5% SDS. For poly A⁺ selection the total RNA sample is heated for 5 minutes at 65°C, adjusted to 0.5
20 M NaCl and subsequently applied to an oligo(dT)-cellulose column. After several washes with an solution containing 10 mM Tris pH 7.0, 1 mM EDTA and 0.1 mM NaCl, the poly A⁺ RNA is collected by elution with 10 mM Tris pH 7.0 and 1 mM EDTA.

Example 2

Preparation and cloning of a cDNA encoding phytase

For the synthesis of the first strand of the cDNA
5 5 µg of poly A⁺ RNA, isolated according to Example 1, is dissolved in 16.5 µl H₂O and the following components are
30 added: 2.5 µl RNasin (30 U/µl), 10 µl of a buffer containing 50 mM Tris-HCl pH 7.6, 6 mM MgCl₂ and 40 mM KCl, 2 µl 1 M KCl, 5 µl 0.1 M DTT, 0.5 µl oligo(dT)₁₂₋₁₈ (2.5 mg/ml), 5 µl
8 mM dNTP-mix, 5µl BSA (1 mg/ml) and 2.5 µl Moloney MLV
35 reverse transcriptase (200 U/µl). The mixture is incubated for 30 minutes at 37°C and the reaction is stopped by

addition of 10 μ l 0.2 M EDTA and 50 μ l H₂O. An extraction is performed using 110 μ l chloroform and after centrifugation for 5 minutes 5 M NH₄Ac and 440 μ l absolute ethanol (-20°C) are added to the supernatant. Precipitation is done in a dry ice/ethanol solution for 30 minutes. After centrifugation (10 minutes at 0°C) the cDNA/mRNA pellet is washed with 70% ice-cold ethanol. The pellet is dried and dissolved in 20 μ l of H₂O.

Isolation of the cDNA encoding phytase is performed with the Polymerase Chain Reaction (PCR) in two fragments. The two fragments are combined, using the BamHI site within the gene to create a full length cDNA. The strategy for the cloning of the phytase cDNA is shown in Figure 1.

Partial sequencing of the phytase gene (Van Gorcom *et al.*, *supra*), reveals the presence of a BamHI site at approximately 800 basepairs from the initiation codon. The nucleotide sequence around this BamHI site, as well as the nucleotide sequence preceding the start codon and the nucleotide sequence after the stop codon of the phytase gene are used to design oligonucleotides for the PCR.

The polymerase chain reaction is performed according to the supplier of Tag-polymerase (Cetus) using 1.5 μ l of the solution containing the reaction product of the first strand synthesis and 0.5 μ g of each of the oligonucleotides. Amplification is performed in a DNA amplifier of Perkin Elmer/Cetus. After 25 cycles of 2 minutes at 94°C, 2 minutes at 55°C, and 3 minutes at 72°C the reaction mixture is deproteinized by subsequent phenol and chloroform extractions. The DNA is precipitated, redissolved in a buffer containing 10 mM Tris, pH 7 and 0.1 mM EDTA and subsequently digested with suitable restriction enzymes.

For the amplification of the fragment encoding the N-terminal part of the protein, the two following oligonucleotides are used:

Oligo 1: 5' GGGTAGAATTCAAAAATGGGCGTCTCTGCTGTTCTA 3'
Oligo 2: 5' AGTGACGAATTCGTGCTGGTGGAGATGGTGTTCG 3'

The amplified fragment is digested with EcoRI and cloned into the EcoRI site of pTZ18R (purchased from Pharmacia). Restriction site mapping and nucleotide sequencing demonstrate the authenticity of the fragment. The resulting plasmid is named pGB925.

For the amplification of the second fragment, the following two oligonucleotides are used:

Oligo 3: 5' GAGCACCAAGCTGAAGGATCC 3'

Oligo 4: 5' AAAGTGCAGGCGTTGAGTGTGATTGTTTAAAGGG 3'

The amplified fragment is digested with BamHI and PstI and subsequently cloned into pTZ18R, which has been digested with BamHI and PstI. Restriction site mapping and nucleotide sequencing show that the correct fragment is isolated. The resulting plasmid is named pGB926.

In order to isolate a full length cDNA, pGB925 is digested with EcoRI and BamHI and the fragment containing the phytase encoding DNA is isolated. This fragment is cloned into plasmid pGB926, which has been digested with EcoRI and BamHI, resulting in plasmid pGB927. Plasmid pGB927 contains a full length cDNA encoding phytase, with an approximate size of 1.8 kbp. The sequence of the cDNA region encoding the phytase protein and the derived amino acid sequence of the phytase protein are depicted in Figure 2.

Example 3

Construction of the binary vector pMOG23.

In this example the construction of the binary vector pMOG23 (in E. coli K-12 strain DH5 α , deposited at the Centraal Bureau voor Schimmel-cultures on January 29, 1990 under accession number CBS 102.90) is described.

The binary vector pMOG23 (Figure 2) is a derivative of vector Bin19 (Bevan, M., supra). To obtain pMOG23, the vector Bin19 is changed in a way not essential for the present invention, using techniques familiar to those skilled in the art of molecular biology.

First, the positions of the left border (LB) and the right border (RB) are switched with reference to the neomycine phosphotransferase gene II (NPTII gene). Secondly, the orientation of the NPTII gene is reversed giving transcription in the direction of LB. Finally, the polylinker of Bin19 is replaced by a polylinker with the following restriction enzyme recognition sites: EcoRI, KpnI, SmaI, BamHI, XbaI, SacI, XhoI, and HindIII.

Example 4

Cloning of the phytase cDNA of *Aspergillus ficuum* in an expression construct for constitutive expression in plants

The phytase gene from *Aspergillus ficuum* is tailored and cloned in an expression construct for constitutive expression downstream of the Cauliflower Mosaic Virus 35S promoter. The expression construct also contains the coding information for a signal peptide sequence of plant origin.

The phytase cDNA is cloned into the expression construct as present on plasmid pMOG29 (described under a)). Subsequently the entire construct is introduced into the binary vector pMOG23 and transferred to *Agrobacterium tumefaciens* strain LBA4404.

a) Construction of expression vector pMOG29

The expression construct of ROK1 (Baulcombe *et al.*, (1986) Nature 321, 446) is cloned as an EcoRI/HindIII fragment into pUC18. This construct contains the Cauliflower Mosaic Virus (CaMV) 35S promoter on an EcoRI/BamHI fragment and the nopaline synthase (nos) transcription terminator on a BamHI/HindIII fragment. The promoter fragment consists of the sequence from -800 to +1 of the CaMV 35S promoter. Position +1, which is included, is the transcription initiation site (Guilley *et al.*, *supra*). The sequence upstream of the NcoI site at position -512 is deleted and this site is changed into an EcoRI site. This is done by cutting the expression construct present in pUC18 with NcoI,

filling in the single-stranded ends with Klenow polymerase and ligation of an EcoRI linker. The resulting plasmid is cut with EcoRI, resulting in the deletion of the EcoRI fragment carrying the sequences of the 35S promoter upstream of the original NcoI site. The BamHI/HindIII fragment, containing the nos terminator is replaced by a synthetic DNA fragment (oligonucleotide duplex A, Figure 4) containing the leader sequence of RNA4 of Alfalfa Mosaic Virus (ALMV) (Brederode *et al.*, *supra*). This is done by cleavage with BamHI, followed by cleavage with HindIII and ligation of the synthetic DNA fragment. The BamHI site and three upstream nucleotides are deleted by site-directed mutagenesis. In the resulting plasmid, the BamHI/HindIII fragment containing the nos terminator sequence is reintroduced. The gene encoding β -glucuronidase (originating from plasmid pRAJ 275; Jefferson, R.A. (1987) *Plant Mol. Biol. Reporter* 5, 387) was ligated in as an NcoI/BamHI fragment, resulting in plasmid pMOG14. From the literature it is known that duplication of the sequence between -343 and -90 increases the activity of the 35S promoter (Kay, R., Chan, A., Dayly, M. & McPherson, J. (1987) *Science* 236, 1299). To obtain a promoter fragment with a double, so-called enhancer sequence, the following steps, known to those skilled in the art, are carried out. From plasmid pMOG14, the enhancer fragment is isolated on an AccI/EcoRI fragment and subsequently blunt-ended with Klenow polymerase. The obtained fragment is introduced in pMOG14 cut with EcoRI and blunt-ended, in such a way that the border between the blunt-ended EcoRI and AccI sites generate a new EcoRI site. The resulting plasmid (pMOG18) contains the 35S promoter with a double enhancer sequence, the leader sequence of RNA4 from ALMV and the nos terminator in an expression construct still present on an EcoRI/HindIII fragment. Finally, the NcoI/BamHI fragment encoding β -glucuronidase is replaced with the synthetic DNA fragment B (Figure 4), derived from the PROB12 cDNA (Cornelissen, B.J.C., Hooft van Huijsduijnen, R.A.M. & Bol, J.F. (1986)

Nature 321, 531). This fragment B encodes the PR-protein PR-S signal peptide sequence from tobacco Samsun NN. An SphI site is created in the signal peptide encoding DNA sequence by changing one nucleotide. This change does not alter the amino acid sequence of the encoded PR-S signal peptide. The resulting plasmid is called pMOG29 (Figure 5).

b) Cloning of the phytase gene from Aspergillus ficuum in the binary vector

Oligonucleotide duplex C (Figure 4) is cloned into plasmid pMOG29, digested with SphI and BamHI, resulting in plasmid pMOG407. The oligonucleotide duplex contains the coding information for the final 2 amino acids of the signal peptide of PR-S, followed by the first 6 amino acids of mature phytase.

The plasmid pGB927, which contains the full length phytase cDNA, is digested with XhoI (partially) and PstI. The XhoI/PstI fragment comprising the DNA sequences encoding mature phytase from amino acid 6 onward is cloned into plasmid pMOG407 linearized with XhoI and PstI, resulting in plasmid pMOG417. The entire construct, containing the chimaeric phytase gene, is inserted as an EcoRI/HindIII fragment into the binary vector pMOG23 linearized with EcoRI and HindIII. The resulting binary plasmid pMOG413 is mobilized, in a triparental mating with the E. coli K-12 strain RK2013 (containing plasmid pRK2013) (Ditta, G., Stanfield, S., Corbin, D. & Helinski, D.R. (1980) Proc. Natl. Acad. Sci. USA 77, 7347), into Agrobacterium tumefaciens strain LBA4404 that contains a plasmid with the virulence genes necessary for T-DNA transfer to the plant.

Example 5

Transient expression of chimaeric phytase gene in tobacco protoplasts

Protoplasts of tobacco are transformed with plasmid DNA carrying the chimaeric phytase gene under regulation of

the constitutive CaMV 35S promoter. After 72 hours treated protoplasts are assayed for transient expression of the introduced phytase gene using the phytase activity assay.

Protoplasts are prepared from axenically grown 1-2 months old tobacco plants (Nicotiana tabacum SR1). The entire procedure is described by Rodenburg, K.W., DeGroot, M.J.A., Schilperoort, R.A. & Hooykaas, P.J.J. ((1989) Plant Mol. Biol. 13, 711). For transformation a number of 5×10^5 protoplasts is electroporated with 40 μ g DNA of plasmid pMOG417). After electroporation protoplasts are resuspended in 3 ml of K3G medium. For the phytase activity assay protoplasts are pelleted and the 3 ml of supernatant is dialyzed overnight against an excess of water. The dialysate is freeze-dried and resuspended in 300 μ l 25 mM sodium-acetate pH 5.5. The assay is then carried out as described in detail in Example 10, with the only exception that instead of the 250 mM Glycine HCl buffer pH 2.5, a 25 mM sodium acetate buffer pH 5.5 is used.

In these experiments, one phytase unit (PTU) is defined as 1 μ mol of phosphate released from 1.5 mM sodium phytate solution per minute at 37°C at pH 5.5.

In untreated protoplasts no detectable activity is found. Protoplasts electroporated with plasmid pMOG417 show an activity of 0.26 PTU per mg protein in the supernatant.

Example 6

Stable expression of a chimaeric phytase gene in tobacco plants under the control of the CaMV 35S promoter

Tobacco is transformed by cocultivation of plant tissue with Agrobacterium tumefaciens strain LBA4404 containing the binary vector pMOG413 with the chimaeric phytase gene under regulation of the CaMV 35S promoter. Transformation is carried out using cocultivation of tobacco (Nicotiana tabacum SRI) leaf discs according to Horsch et al., supra. Transgenic plants are regenerated from shoots that grow on selection medium (100 mg/l) kanamycin), rooted

and transferred to soil. Young plants are assayed for NPTII-activity (kanamycin resistance), grown to maturity and allowed to self-pollenate and set seed.

5 For phytase activity assays of the leaves of the transgenic plants, a segment of approx. 5 mm in diameter from a young leaf is taken from each plant, and homogenized in 300 μ l 25 mM sodium-acetate buffer pH 5.5. Subsequently, phytase assays were carried out as described for the transient assays. In 32 independently transformed tobacco
10 plants tested, a maximum activity was observed of 2 PTU/mg total soluble protein in the extracts. This corresponds to 1.7% of total soluble protein. In the seeds of these transformed tobacco plants, a maximum phytase expression level of 0.4% of the total soluble seed protein was
15 observed. No phytase activity could be detected in untransformed plants.

Two transgenic plant lines, 413.25 and 413.32, were selected on the basis of their high expression levels of phytase.

20

Example 7

Cloning of the phytase cDNA of Aspergillus ficuum in a seed-specific expression construct

25 An expression construct is constructed in such a way that seed-specific expression is obtained, using sequences of the Brassica napus 12S storage protein gene cruciferin (cruA; Ryan et al., supra). These sequences may be replaced by those from similar seed-specific genes to achieve the same goal as is the objective of this invention.

30 The phytase cDNA is cloned into the expression construct. Finally, the entire construct is introduced into Agrobacterium tumefaciens, which is used for transformation.

For all E. coli transformations in this example, E. coli K-12 strain DH5 α is used.

35

a) Construction of the expression construct

For the construction of the expression construct for seed-specific expression, the promoter and terminator sequences from the cruciferin A (cruA) gene of Brassica napus cv. Jet Neuf are synthesized using PCR technology with isolated genomic DNA (Mettler, I.J. (1987) Plant Mol. Biol. Rep. 5, 346) as a template. This gene shows seed-specific expression and its coding and flanking sequences have been determined (Ryan et al., supra).

Two sets of oligonucleotides are synthesized. One to allow amplification of the cruA 5' flanking region and part of the signal peptide encoding sequence as an EcoRI/NcoI fragment:

5' GTTCGGAATTCGGGTTCCGG 3' and 5' AACTGTTGAGCTGTAGAGCC 3'.
The other for amplification of the 3' flanking sequence as a BglII/HindIII fragment:

5' CTTAAGATCTTACCCAGTGA 3' and 5' CGGAGAAGCTTGCATCTCGT 3'.

The oligo's are designed to contain suitable restriction sites at their termini to allow direct assembly of the expression construct after digestion of the fragments with the restriction enzymes.

The 5' fragment of the cruA gene, that includes 54 nucleotides of the sequence encoding the signal peptide is cloned into vector pMOG445 (Oligonucleotide duplex E (Figure 4) cloned into vector pUC18, linearized with SstI and EcoRI), cut with EcoRI and NcoI, resulting in vector pMOG424. The synthetic oligonucleotide duplex D (Figure 4), comprising the final 5 coding triplets for the signal sequence of Brassica napus cruciferin, the sequence encoding amino acids 1-6 of mature phytase and a multiple cloning site, is cloned in vector pMOG424 cut with NcoI and HindIII. The resulting vector is called pMOG425. The 3' cruA PCR fragment is cloned as a BglII/HindIII fragment into pMOG425 digested with BglII and HindIII, resulting in pMOG426.

b) Cloning of the phytase gene from Aspergillus ficuum in the binary vector

Plasmid pGB927, which contains the full-length coding sequence for Aspergillus ficuum phytase, is digested with XhoI (partially) and with PstI. The XhoI/PstI fragment comprising the DNA sequences encoding mature phytase from amino acid 6 onward is cloned in vector pMOG426, cut with XhoI and PstI. From the resulting vector pMOG428, the entire construct, containing the chimeric phytase gene, is inserted as an EcoRI/HindIII fragment in the binary vector pMOG23 linearized with EcoRI and HindIII. The resulting binary vector pMOG429 is mobilized, in a triparental mating with the E. coli K-12 strain RK2013 (containing plasmid pRK2013) (Ditta et al., supra), into Agrobacterium strain LBA4404 (Hoekema et al., 1983, supra) that contains a plasmid with the virulence genes necessary for T-DNA transfer to the plant.

Example 8

Stable seed-specific expression of phytase in tobacco seeds under the control of a cruciferin promoter

Agrobacterium strain LBA4404, containing the binary vector pMOG429 with the phytase cDNA under the control of the cruciferin promoter, is used for transformation experiments. Transformation of tobacco (Nicotiana tabacum SR1) is carried out using cocultivation of leaf discs according to the procedure of Horsch et al., supra. Transgenic plants are regenerated from shoots that grow on selection medium (100 mg/l kanamycin). Young plants are assayed for NPTII-activity (kanamycin resistance), grown to maturity and allowed to self-pollenate and set seed. Seeds from individual transformants are pooled and part of the seed sample is assayed for the presence of phytase. From clones with the highest expression levels, compared to untransformed control seeds, the remaining seeds are germinated on kanamycin (200 mg/L). From data on the resulting S2 seeds, seeds homozygous for NPTII (hence also for phytase) are selected and used for mass propagation of

plants capable of producing the highest amounts of phytase. These can then be used, e.g. for digestion experiments.

To determine the phytase activity found in the transgenic seeds, about 50 mg seed is taken and homogenized with a pestle in an ice-cold mortar in 1 ml 25 mM sodium-acetate buffer pH 5.5. After centrifugation, the supernatant is assayed as described for the transient assays. In 55 independently transformed tobacco plants, a maximum phytase expression level of 0.15% of the total soluble seed protein was observed. Phytase activity was not detected in stems, roots and leaves of the transgenic plants. No phytase activity could be detected in untransformed plants.

Example 9

Transformation of rapeseed

In this example, the transformation of rapeseed by co-cultivation of plant tissue with Agrobacterium tumefaciens, containing a binary vector with the chimeric phytase gene is described. Transgenic plants may be selected on antibiotic resistance. The transgenic plants may be assayed for phytase activity. High expressors may be analyzed more thoroughly and used in further experiments.

The same chimeric phytase construct in a binary vector (pMOG429) is mobilized into Agrobacterium tumefaciens strain LBA4404, in a like manner as described in Example 7. This strain may be used to transform rapeseed (Brassica napus cv. Westar). To this aim, surface-sterilized stem segments taken from 5 to 6 week-old plants, just before flowering, are preconditioned for 24 h on MS medium (Fry et al. (1987) Plant Cell Reports 6, 321) with 1 mg/l BAP and then co-cultivated for 48 h with Agrobacterium on fresh plates with the same medium. Transgenic plantlets were regenerated from shoots that grow on selection medium (500 mg/l carbenicilline, 40 mg/l paromomycin) and further analyzed as described in Example 8 for tobacco.

Example 10

Phytase activity assay

An amount of transgenic plant material was ground which in total contain approximately 0.25 PTU. (PTU = Phytase units. One unit of phytase activity is defined as that amount of enzyme which liberates inorganic phosphorus from 1.5 mM sodium phytate at the rate of 1 $\mu\text{mol/min}$ at 37°C and at pH 2.5). Alternatively, this amount of phytase may be extracted from the plant material.

The ground plant material was incubated in a total volume of 50 ml of a 250 mM glycine/HCl buffer pH 2.5 containing 0.86 g sodium phytate $\cdot 11 \text{ H}_2\text{O}$. Although Aspergillus phytase expresses a pH optimum at 2.5 as well as at 5.5, the lower pH is chosen to exclude plant phytase activity.

The resulting mixture is incubated for 15 and 60 minutes at 37°C. The reaction is stopped by the addition of 5 ml from the incubate into 5 ml of 10% TCA (trichloroacetic acid). Thereafter, 10 ml of indicator reagent (3.66 g of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ in 50 ml of ammonium molybdate solution (2.5 g $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$ and 8 ml conc. H_2SO_4 , diluted up to 250 ml with demiwater) is added to the stopped enzyme solution. The intensity of the blue color is measured spectrophotometrically at 700 nm.

The inorganic phosphate content present at $T = 0$ serves as a blank.

The measurements are indicative of the quantity of phosphate released in relation to a calibration curve of phosphate in the range of 0-1 mM.

Example 11

Incubation of ground *Nicotiana tabacum* plant material with feedstuffs

In a typical experiment, 0.25 g of solvent extracted soybean meal is incubated with a quantity of ground *Nicotiana tabacum* plant material containing approximately

0.25 PTU as described above, except for the addition of sodium phytate. In this case, the added incubation agent consists of a mixture of 410 ml buffer and 90 ml of demiwater.

5 The liberation of phosphate from phytate in solvent extracted soybean meal is depicted in Figure 6. Without added ground plant material, no activity is observed.

 In a virtually identical experiment, similar results are obtained using maize gluten feed as a substrate. Results using transgenic seeds are shown in Figure 6.

10 No activity is observed in the absence of ground plant material or when ground plant material are added which do not contain phytase activity.

15 Example 12

In vitro testing of transgenic plant material containing
 phytase under conditions simulating the digestive
 tract of poultry

20 To assess the effectivity of phytase produced in transgenic tobacco plant material, the activity of phytase from Aspergillus is determined in a model simulating the conditions found in the digestive tract in poultry.

25 A standard poultry feed sample is first incubated at 1 g/15 ml demi water for 60 minutes at 39°C to simulate the conditions in the crop of the animals. Subsequently, 5 ml of a pepsin solution (Merck: 5.28 g/l, pH 3.0 - adjusted with HCl) is added, the pH adjusted with HCl to pH 3.0, and the incubation is continued for a further 90 minutes at the same temperature to simulate the conditions in the stomach.

30 During the incubation period, samples were taken in order to determine the amount of phosphate released from the phytate present in the feed.

35 The action of fungal phytase is apparent from Figure 7. Increasing the phytase dosage from 250 to 1000 PTU/kg feed results in an increased release of phosphate from the feed sample.

When a sample of transgenic tobacco plant material, either seed or leaf (lines 413.25 and 413.32; after grinding in a mortar), is added in place of the fungal phytase, a similar increased phosphate release is observed (Figure 8).

5 Control tobacco plant material, which did not contain phytase, was also tested. No phosphate release was observed as compared to the blank control.

Comparison of the results with 50 g transgenic tobacco seed/kg feed with those obtained with 500 and 750
10 PTU/kg feed indicates that 1 g tobacco seed equals approximately 12 PTU in this in vitro poultry digestion model. A sample comparison using leaf material indicates that 1 g (fresh weight) of tobacco leaf material contains approximately 25 PTU.

15

Example 13

Animal testing

Trials are carried out with broilers to show the efficacy of phytase expressed in plant seeds, as well as the
20 absence of any negative effect of seeds from tobacco on zootechnical results.

Both phytase-expressing and control tobacco seed are harvested. Seeds were ground in 100 gram portions with a sieve (Retch-mill ZM1) having pores of 500 μ m, taking care
25 to keep the seeds cooled.

One day old male broiler chicks (Hybro) are housed in two tier battery cages (0.45 m²). The ambient temperature is 32°C during the first two days and is decreased by 4°C in the first week. Every following week, the temperature is
30 decreased by 2°C. Broilers are reared in a one hour light and three hours dark regime.

The birds are vaccinated against New Castle Disease at one day of age using Clone 30 vaccine. During the experiments, the broilers are fed the experimental diets all
35 mash and ad libitum. Growth and feed/gain ratios are measured during the experimental periods. Apparent

availability of total phosphorus is measured in a three day period, during which feed consumption is measured as dry matter intake and excreta are collected quantitatively.

5 Apparent availability of phosphorus is defined as the difference between intake of phosphorus and excretion of phosphorus with the excreta.

The following control diets without addition of phytase are used:

10	Diets	Ca (%)	total P (%)	phytate P (%)
	1	0.60	0.45	0.30
	2	0.75	0.60	0.30
	3	0.90	0.75	0.30

15 No graded feed phosphate is added to diet 1 (basal diet). Calcium and phosphorus from a mixture of anhydrous dicalcium phosphate and monoammonium phosphate (ratio 5:1) are supplemented to diets 2 and 3. All experimental diets are obtained by additions to the basal diet (see Table 1).

20 Experimental diet 4 contains microbial phytase at a concentration of 400 PTU/kg feed, prepared as described by Van Gorcom et al., supra.

25 Experimental diet 5 is like diet 4 but ground seeds of non-transgenic tobacco are added to the feed mixture to achieve a final ratio of 3 kg/90 kg feed.

Experimental diet 6 is also like diet 4 but 3 kg ground seeds of transgenic tobacco (line 413.25) are added to a mixture of 90 kg feed to obtain a final concentration of 400 PTU/kg feed.

30 The experiment is carried out with 176 broilers in 16 battery cages (11 per battery cage) until the age of 24 days. The treatments (diets) are repeated twice and are assigned randomly to the cages within each tier.

35 The availability of phosphorus is measured from 21-24 days of age.

5 The results with regard to phosphorous availability and growth of the animals supplied with diets 4, 5 and 6 each show the positive effect of the addition of phytase (Table 2). A comparison of diets 4, 5 and 6 also demonstrates that the inclusion of tobacco seeds in feed is compatible with the action of microbial phytase in the gastro-intestinal tract of farm animals such as broilers and shows no negative effect on the zootechnical results.

5 While the present invention has been described with reference to specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without parting from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, plant, seed, process, process step or steps to the object, spirit and scope of the
10 invention. All such modifications are intended to be within the scope of the claims appended hereto.

Table 1 Composition of basal diet in experiments with broilers

5	Ingredients	Contents (g/kg)
	Yellow maize	280.0
10	Sorghum (low tannin)	200.0
	Sunflower seed meal (solv. extr.)	80.0
	Soya bean meal (solvent extr., 48.8% protein)	350.0
	Soya bean oil	58.5
	Vitamins*	5.0
15	Minerals*	15.0
	Limestone	1.0
	Synth. methionine	1.0
	Cr ₂ O ₃	0.5
		1001.0
20	ME (MJ/kg)	13.1
	Lysine	12.9
	Methionine + cystine	9.1
	Calcium	6.0 (6.0 - 6.6)**
25	Total phosphorus	4.5 (4.7 - 4.7)**
	Organic phytic phosphorus	3.0 (3.1 - 3.1)**

* Amount supplied per kg diet: 12000 IU vitamin A; 2000 IU vitamin D₃; 5 IU vitamin E; 1.5 mg vitamin K₃; 1 mg thiamine; 5 mg riboflavin; 1 mg pyridoxine; 30 mg nicotinic acid, 7.5 mg D-pantothenic acid; 0.015 mg vitamin B₁₂; 0.5 mg folic acid; 350 mg choline chloride; 75 mg ethoxyquin; 9.5 g CaCO₃; 2.5 g NaCl; 0.26 g FeSO₄; 0.24 g MnSO₄; 45 mg CuSO₄; 60 mg ZnSO₄; 105 mg KI mixture.

** () Analyzed for experiments 1 and 2 respectively.

Table 2
The Effect of Phytase on the Apparent
Availability of Total P and Ca, the P Content in
Manure and the Performance of Broilers

10	Diets	Ca/P (g/kg)	Added phytase (units/kg)	Availability (%)		Amount of P in manure (g) per kg dm feed intake	Growth 0-24 d (g)
				21-24 d	P Ca		
15	1	6/4.5	0	49.8	47.2	2.7	338
	2	7.5/6	0	45.6	48.9	3.8	592
	3	9/7.5	0	44.6	46.9	4.9	683
15	4	as 1	400	60.5	58.6	2.1	620
	5	as 1	0	48.5	48.0	2.7	340
	6	as 1	400	60.2	59.3	2.1	615

CLAIMS

1. A method for the production of transgenic plants or plant organs containing an enhanced amount of a phytase characterized in that a plant host is transformed with an expression construct which contains a DNA sequence encoding the phytase operably linked to regulatory sequences capable of directing the expression of the phytase in the plant host and growing the transformed plant under conditions conducive whereby the phytase-encoding DNA sequence is expressed.
2. The method of Claim 1 wherein the expression construct contains a regulatory sequence capable of directing the constitutive expression of the phytase.
3. The method of Claim 1 wherein the expression construct is capable of directing the tissue-specific expression of the phytase.
4. The method of Claim 1 wherein the phytase-encoding DNA sequence is obtained from a microbial source.
5. The method of Claim 4 wherein the phytase-encoding DNA sequence is obtained from a filamentous fungus source.
6. The method of Claim 5 wherein the phytase-encoding DNA sequence is obtained from an Aspergillus source.
7. The method of Claim 6 wherein the phytase-encoding DNA sequence is obtained from an Aspergillus source selected from the group consisting of A. ficuum, A. niger, A. awamori and A. nidulans.
8. An expression construct characterized in that a DNA sequence encoding a phytase is operably linked to a regulatory sequence capable of directing the constitutive expression of the phytase.

9. A vector characterized in that the vector contains an expression construct according to Claim 8.

10. A bacterial strain characterized in that the strain contains a vector according to Claim 9.

11. An expression construct characterized in that a DNA sequence encoding a phytase is operably linked to a regulatory sequence capable of directing the tissue-specific expression of the phytase.

12. A vector characterized in that the vector contains an expression construct according to Claim 11.

13. A bacterial strain characterized in that the strain contains a vector according to Claim 12.

14. A transgenic plant or plant organ characterized in that the plant or plant organ contains an expression construct which contains a DNA sequence encoding a phytase operably linked to regulatory sequences capable of directing the expression of the phytase in the plant host.

15. A method for the conversion of phytate to inositol and inorganic phosphate characterized in that a phytase produced according to the method of any one of Claims 1 to 7 is applied to a substrate which contains phytate.

16. A feed composition for animals characterized in that the feed contains a phytase produced according to the method of any one of Claims 1 to 7.

17. The feed composition of Claim 16 characterized in that the feed contains a transgenic plant or plant organ containing phytase produced according to the method of any one of Claims 1 to 7.

18. A method for promoting the growth of animals characterized in that an animal is fed a diet which is comprised of a feed according to Claim 16 in an amount effective in promoting the growth of the animal ingesting the feed.

19. A method for the reduction of levels of phytate in animal manure characterized in that an animal is fed a diet which is comprised of a feed according to Claim 16 in an amount effective in converting phytate contained in the feed to inositol and inorganic phosphate.

20. A method for the production of phytase characterized by:

- a) transforming a plant host with an expression construct which contains a DNA sequence encoding the phytase operably linked to regulatory sequences capable of directing the expression of an enhanced amount of phytase in the plant host and growing the transformed plant under conditions conducive whereby the phytase-encoding DNA sequence is expressed in the plant tissue; and
- b) extracting the phytase from the transgenic plant tissue.

21. Use of phytase produced according to the method of any one of Claims 1 to 7 and 21 as an ingredient in food or feedstuffs.




Figure 1

SUBSTITUTE SHEET

FIGURE 2 (Sheet 1 of

1 ATGGGCGTCTCTGCTGTTCTACTTCCTTTGTATCTCCTGTCTGGAGTCACCTCCGGACTG
-23 M G V S A V L L P L Y L L S G V T S G L

61 GCAGTCCCCGCTCGAGAAATCAATCCAGTTGCGATACGGTCGATCAGGGGTATCAATGC
A V P A S R N Q S S C D T V D Q G Y Q C
-1 +1

121 TTCTCCGAGACTTCGCATCTTTGGGGTCAATACGCACCGTTCCTTCTCTCTGCGAAACGAA
18 F S E T S H L W G Q Y A P F F S L A N E

181 TCGGTCATCTCCCCTGAGGTGCCCCGCGGATGCAGAGTCACTTTGCTCAGGTCTCTCTCC
38 S V I S P E V P A G C R V T F A Q V L S

241 CGTCATGGAGCCGCGTATCCGACCGACTCCAAGGGCAAGAAATACTCCGCTCTCATTGAG
58 R H G A R Y P T D S K G K K Y S A L I E

301 GAGATCCAGCAGAACGCGACCACTTTGACGGAAAATATGCCTTCCTGAAGACATACAAC
78 E I Q Q N A T T F D G K Y A F L K T Y N

361 TACAGCTTGGGTGCAGATGACCTGACTCCCTTCGGAGAACAGGAGCTAGTCAACTCCGGC
98 Y S L G A D D L T P F G E Q E L V N S G

421 ATCAAGTTCTACCAGCGGTACGAATCGCTCACAAGGAACATCGTTCCATTTCATCCGATCC
118 I K F Y Q R Y E S L T R N I V P F I R S

481 TCTGGCTCCAGCCGCGTGATCGCCTCCGGCAAGAAATTCATCGAGGGCTTCAGAGCACC
138 S G S S R V I A S G K K F I E G F Q S T

541 AAGCTGAAGGATCCTCGTGCCCGAGCCCGGCCAATCGTCGCCCCAAGATCGACGTGGTCATT
158 K L K D P R A Q P G Q S S P K I D V V I

601 TCCGAGGCCAGCTCATCCAACAACACTCTCGACCCAGGCACCTGCACTGTCTTCCAAGAC
178 S E A S S S N N T L D P G T C T V F E D

661 AGCGAATTGGCCGATACCGTGAAGCCAATTTACCGCCACGTTGCTCCCTCCATTTCGT
198 S E L A D T V E A N F T A T F V P S I R

721 CAACGTCTGGAGAACGACCTGTCCGGTGTGACTCTCAGACACAGAAAGTGACCTACCTC
218 Q R L E N D L S G V T L T D T E V T Y L

781 ATGGACATGTGCTCCTTCGACACCATCTCCACCAGCACCGTCGACACCAAGCTGTCCCCC
238 M D M C S F D T I S T S T V D T K L S P

SUBSTITUTE SHEET

FIGURE 2 (Sheet 2 of 2)

841 TTCTGTGACCTGTTACCCATGACGAATGGATCAACTACGACTACCTCCAGTCCTTGAAA
258 F C D L F T H D E W I N Y D Y L Q S L K

901 AAGTATTACGGCCATGGTGCAGGTAACCCGCTCGGCCCGACCCAGGGCGTCGGCTACGCT
278 K Y Y G H G A G N P L G P T Q G V G Y A

961 AACGAGCTCATCGCCCGTCTGACCCACTCGCCTGTCCACGATGACACCAGTTCCAACCAC
298 N E L I A R L T H S P V H D D T S S N H

1021 ACTTTGGACTCGAGCCCGGCTACCTTTCCCTCAACTCTACTCTCTACGGGACTTTTCG
318 T L D S S P A T F P L N S T L Y A D F S

1081 CATGACAACGGCATCATCTCCATTCTCTTTGCTTTAGGTCTGTACAACGGCACTAAGCCG
338 H D N G I I S I L F A L G L Y N G T K P


1141 CTATCTACCACGACCGTGGAGAATATCACCCAGACAGATGGATTCTCGTCTGCTTGGACG
358 L S T T T V E N I T Q T D G F S S A W T

1201 GTTCCGTTTGCTTCGCGTTTGTACGTCGAGATGATGCAGTGTGAGCCGGAGCAGGAGCCG
378 V P F A S R L Y V E M M Q C Q A E Q E P

1261 CTGGTCCGTGTCTTGGTTAATGATCGCGTTGTCCCGCTGCATGGGTGTCCGGTTGATGCT
398 L V R V L V N D R V V P L H G C P V D A

1321 TTGGGGAGATGTACCCGGGATAGCTTTGTGAGGGGGTTGAGCTTTGCTAGATCTGGGGGT
418 L G R C T R D S F V R G L S F A R S G G

1381 GATTGGGCGGAGTGTTTTGCTTAG
438 D W A E C F A *



Polylinker sequence:

Eco RI	Kpn I	Sma I	Bam HI	Xba I	Hind III
5' GGAATTCTGGTACCTCCCGGAGGATCCATCTAGAGCTCGAGTAAGCTTC 3'					
				Sac I	

Figure 3

SUBSTITUTE SHEET

FIGURE 4 (Sheet 1 of 2)

Oligonucleotide duplex A

5'GGGTTTATTTTAAATTTCTTTCAAAATACCTCCACCATGGGTAACGGATCCA 3'
 3'CCCAAAATAAAAATTAAGAGTTTATGAAGGTGTACCCATGTCCTAGGTTCCA 5'

NcoI BamHI HindIII

[] [] [] [] [] []

Oligonucleotide duplex B

5'CATGAAACTTCCTCAAAAGCTTCCCTTTTATGCTTCTTGTGTTGGCCAAATACCTTGTAGCTGTACGCATGCTCGAG 3'
 3' TTTGAAGGAGTTTTCGAAGGGGAAAAATACGGAAGGAAACAAAACCGGTTATGAAACATCGACAATGCGTACGAGCTCCTAG 5'

SphI XhoI BamHI

[] [] [] [] [] []

Oligonucleotide duplex C

5' CT CTGGCAGTCCCGCCCTCGAGCCCCCTGCAG 3'
 3' GTACGA GACCGTCAGGGCGGAGCTCGGGGGACGTCCTAG 5'

SphI XhoI PstI BamHI

[] [] [] [] [] []

PROB12 Mature phytase
 signal peptide

Oligonucleotide duplex D

Oligonucleotide duplex E

5' AATCAGATCTCCATGGATCGATGAGCT 3'
3' GTCTAGAGGTACCTAGCTAC 5'




Figure 5




FIGURE 6

IN VITRO DIGESTIBILITY MODEL

ASPERGILLUS PHYTASE DOSE-RESPONSE RELATION




Figure 7

IN VITRO DIGESTIBILITY MODEL

ASPERGILLUS & TOBACCO SEED PHYTASE





Figure 8

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM SECOND SHEET)		
Category *	Citation of Document, " with indication, where appropriate, of the relevant passages	Relevant to Claim No.
X	Patent Abstracts of Japan, volume 9, no. 17 24 January 1985 (C-262)(1740), & JP, A, 59166049 (NIPPON SHINYAKU K.K.) 19 September 1984 see the abstract ---	15,16
X	EP, A, 0287152 (GIST BROCADES) 19 October 1988 see the whole document ---	19
Y	The Journal of Cellular Biochemistry, Sup- plement 12C, 17th Annual Meetings, 28 February-10 April 1988, Alan R. Liss, Inc., (New York, US), A.A. Christen et al.: "Cloning of the phytase gene from germinating soybeans", page 190, see abstract L 402 ---	1-3,8-14,20
Y	WO, A, 87/00865 (CALGENE) 12 February 1987 see page 4, lines 10-16; page 9, line 32 - page 10, line 4 ---	1-3,8-14,20
X	US, A, 4859486 (DOUGLASS) 22 August 1989 see column 5, lines 15-23 ---	21
P,X	EP, A, 0380343 (ALKO) 1 August 1990 see the whole document ---	15
E	EP, A, 0420358 (GIST-BROCADES) 3 April 1991 see the whole document ---	8-10,15-19, 21
A	WO, A, 90/01551 (ROGERS) 22 February 1990 see page 12, lines 1-10; page 12, lines 21-32 ---	20
X	EP, A, 0346909 (AALTONEN) 20 December 1989 see page 3, lines 8-10 ---	21
X	US, A, 3297548 (WARE) 10 January 1967 see the whole document, in particular column 1, lines 25-33 -----	15,16,18

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols indicate all) *		
According to International Classification (IPC) or to both National Classification and IPC		
IPC ⁵ : C 12 N 15/82; 15/55, 9/16, 1/21, A 23 K 1/165, A 01 H 5/00		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
IPC ⁵	C 12 N, A 23 K, A 01 H	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	Journal of Agricultural & Food Chemistry, volume 36, no. 6, American Chemical Society, (Washington, DC, US), Y.W. Han et al.: "Phytate hydrolysis in soybean and cottonseed meals by <i>Aspergillus ficuum</i> phytase", pages 259-262 see the whole document	15,16,21
X	EP, A, 0321004 (DORR-OLIVER) 21 June 1989 see the whole document	15
X	EP, A, 0179441 (SIREN) 30 April 1986 see the whole document	16,21
X	Journal of Agricultural & Food Chemistry, volume 36, no. 2, American Chemical Society, (Washington, DC, US), Y.W. Han: "Removal of phytic acid from soybean and cottonseed meals", pages 1181-1183 see page 1181, left-hand column	16,18,21
--- ./.		
<p>* Special categories of cited documents: ¹⁴</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
26th June 1991	29.08.91	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	F.W. HECK 	

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

NL 9100047
SA 46143

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 20/08/91. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0321004	21-06-89	NL-A- 8702735	16-06-89
		JP-A- 1281048	13-11-89
		US-A- 4914029	03-04-90
EP-A- 0179441	30-04-86	AU-B- 589378	12-10-89
		AU-A- 4893085	01-05-86
		AU-B- 582594	06-04-89
		AU-A- 4893185	01-05-86
		AU-B- 596080	26-04-90
		AU-A- 4893285	01-05-86
		AU-B- 598727	05-07-90
		AU-A- 4893385	17-07-86
		BE-A- 903497	17-02-86
		BE-A- 903498	17-02-86
		CH-A- 666812	31-08-88
		CH-A- 668597	13-01-89
		DE-A- 3537542	15-05-86
		DE-A- 3537550	15-05-86
		DE-A- 3537563	15-05-86
		DE-A- 3537569	22-05-86
		DE-A- 3565464	17-11-88
		EP-A,B 0179439	30-04-86
		EP-A,B 0179440	30-04-86
		EP-A,B 0179442	30-04-86
		FR-A,B 2571967	25-04-86
		FR-A,B 2573080	16-05-86
		GB-A,B 2167298	29-05-86
		GB-A- 2169602	16-07-86
		GB-A,B 2167283	29-05-86
		GB-A,B 2167420	29-05-86
		JP-A- 61171422	02-08-86
		JP-A- 61171491	02-08-86
		JP-A- 61181347	14-08-86
		JP-A- 61171789	02-08-86
		SE-A- 8405295	24-04-86
		SE-A- 8503164	24-04-86
		SE-A- 8503165	24-04-86
		SE-A- 8504967	24-04-86
		SE-A- 8504968	24-04-86

EPO FORM 10079

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.

NL 9100047
SA 46143

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 20/08/91. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0179441		SE-A- 8504969	24-04-86
		SE-C- 458121	22-06-89
		SE-A- 8504970	24-04-86
		US-A- 4794014	27-12-88
		US-A- 4797390	10-01-89
		US-A- 4793945	27-12-88
		US-A- 4851560	25-07-89
		US-A- 5003098	26-03-91
		US-A- 5023248	11-06-91
		US-A- 4735936	05-04-88
		US-A- 4734283	29-03-88
		US-A- 4777134	11-10-88
		US-A- 4735902	05-04-88
EP-A- 0287152	19-10-88	JP-A- 63269990	08-11-88
WO-A- 8700865	12-02-87	US-A- 4956282	11-09-90
		AU-B- 603063	08-11-90
		AU-A- 6196686	05-03-87
		EP-A- 0233915	02-09-87
		JP-T- 63500425	18-02-88
US-A- 4859486	22-08-89	None	
EP-A- 0380343	01-08-90	WO-A- 9008476	09-08-90
EP-A- 0420358	03-04-91	AU-A- 6501190	28-04-91
		WO-A- 9105053	18-04-91
WO-A- 9001551	22-02-90	AU-A- 4037289	05-03-90
		EP-A- 0428572	29-05-91
EP-A- 0346909	20-12-89	JP-A- 2042945	13-02-90
US-A- 3297548		DE-B- 1300488	
		GB-A- 1064304	
		NL-A- 6509747	31-01-66

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82